

We are proud to present the first article within the MARBEFES project, among others.

This is an article in **Ecological Indicators**, Elsevier !

Cite: Paolo Magni, Seyed Ehsan Vesal, Jacopo Giampaolotti, Serena Como, Maria Flavia Gravina. Joint use of biological traits, diversity and biotic indices to assess the ecological quality status of a Mediterranean transitional system, *Ecological Indicators*, Vol. 147, 2023, 109939, ISSN 1470-160X, <https://doi.org/10.1016/j.ecolind.2023.109939> (<https://www.sciencedirect.com/science/article/pii/S1470160X2300081X>)



Joint use of biological traits, diversity and biotic indices to assess the ecological quality status of a Mediterranean transitional system

Paolo Magni ^{a,b,*}, Seyed Ehsan Vesal ^a, Jacopo Giampaolotti ^a, Serena Como ^c, Maria Flavia Gravina ^{a,d,e}

^a Consiglio Nazionale delle Ricerche, Istituto per lo studio degli impatti Antrici e Sostenibilità in ambiente marino (CNR-IAS), Loc. S. Marini, Torregreca, Oristano 09170, Italy

^b Centro Marino Internazionale (CMI), Loc. S. Marini, Torregreca, Oristano 09170, Italy

^c Consiglio Nazionale delle Ricerche, Istituto di Biofisica (CNR-IBF), Pisa 56124, Italy

^d Consorzio Nazionale Interuniversitario per le Scienze del Mare (CoNISMa), Piazzale Flaminio 9, Rome 00196, Italy

^e Università di Roma "Tor Vergata", Dipartimento di Biologia, Via della Ricerca Scientifica, 1, Rome 00133, Italy

ARTICLE INFO

Keywords:
Benthic macroinvertebrates
Biotic indices
Benthic indicators
Biological trait analysis
Functional diversity
Water Framework Directive (WFD)
Coastal lagoons

ABSTRACT

In the present study, we tested the effectiveness of biological trait analysis (BTA) as an ecological indicator in support of the assessment of the Ecological Quality Status (EQS) of a Mediterranean transitional system by commonly used macrobenthic assemblage diversity (e.g. H' and Margalef) and biotic (BENTIX, M-AMBI and M-BAMBI) indices. Ten biological traits describing morphological, behavioral and life history characteristics were considered for the analysis of the functional structure of macrobenthos. As a test study, a historical dataset obtained from the Cabras lagoon (Gardinia, Italy) comprising a riverine site, the lagoonal main body, a confined pond, a seaward creek and a marine channel connected to the Gulf of Oristano was analyzed. Our results showed clear spatial changes in macrobenthic density, number of species and diversity indices along the riverine-lagoonal gradient, which were well reflected in the EQS values based on M-AMBI and to a lesser extent on M-BAMBI. The abundance-based functional diversity was significantly related to the structure and taxonomic diversity of the benthic community. Similarly, the Community-level Weighted Mean test computed with abundance data, rather than with biomass data, supported major evidence for differences in the ecological conditions of the different study sites. In particular, all riverine, lagoonal and confined stations had poor or bad EQS, indicating unstable and warning conditions in this Mediterranean transitional water body. Overall, the BTA approach appeared to be a very promising method in reporting the expected environmental and biotic variability in the Cabras lagoon system and in supporting the EQS assessment by M-AMBI. We conclude that an in-depth knowledge of the benthic communities in terms of both taxonomic and functional aspects is an indispensable basic tool for understanding the variability of the brackish-water systems and for properly assessing their environmental quality.

[authority]20[/authority]

Odpowiedzialny za treść:	Paulina
Wytworzył:	IO PAN
Data wytworzenia:	20.02.2023
Opublikował w BIP:	Paulina Pakszys

Data opublikowania:	21.02.2023 14:52
Ostatnio zaktualizował:	Paulina Pakszys
Data ostatniej aktualizacji:	26.05.2023 16:32
Liczba wyświetleń:	1904